Erratum to “Effect of increasing oil from distillers grains or corn oil on lactation performance” (J. Dairy Sci. 88:2820–2827)

C. Leonardi, S. Bertics, and L. E. Armentano

Table 4 (page 2826) contained incorrect data for milk fatty acids C10:0, C12:0, and C14:0. In the original published table, numerical data for C10:0 were inadvertently omitted, resulting in an upward shift of data rows. Thus, C12:0 data were mistakenly reported as C10:0 data, C14:0 data as C12:0 data, and sum of C4:0 through C12:0 were reported as C14:0 data. Corrected data for C10:0, C12:0, and C14:0 are reported below.

The authors regret the error.

Table 4. Treatment effect on milk fatty acid profile

Profile	Diet ¹					SEM	Statistical contrast ($P < F$) ²			
	0%	5%	10%	15%	OIL		Linear	Quadratic	OIL vs. 0%	OIL vs. 15%
	g/100 g of fatty acids									
C _{10:0}	3.18	3.07	2.98	2.83	2.73	0.16	0.0001	0.74	<0.0001	0.28
C _{12:0}	3.57	3.42	3.31	3.17	3.02	0.15	<0.0001	0.94	<0.0001	0.11
C _{14:0}	11.0	10.7	10.3	10.0	10.0	0.3	<0.0001	0.86	<0.0001	0.88

¹Diets: 0, 5, 10, or 15% DGS = diets containing distillers dried grains with solubles (DGS, DM basis); OIL = diet containing corn oil but no DGS; the amount of corn oil added was such that OIL and 15% DGS had similar total fatty acid concentration.

REFERENCES

Leonardi, C., S. Bertics, and L. E. Armentano. 2005. Effect of increasing oil from distillers grains or corn oil on lactation performance. J. Dairy Sci. 88(8):2820–2827.