Short communication: Cheese supplemented with *Thymus algeriensis* oil, a potential natural food preservative

Danka Bukvicki,^{*1} Abdulhmid Giweli,[†] Dejan Stojkovic,[‡] Ljubodrag Vujisic,[§] Vele Tesevic,[§] Milos Nikolic,[‡] Marina Sokovic,[‡] and Petar D. Marin^{*}

^{*}University of Belgrade - Faculty of Biology, Institute of Botany and Botanical Garden "Jevremovac," 11000 Belgrade, Serbia

[†]Department of Botany, Faculty of Science, University of Al-Gabel Al-Garbe, Zintan, Libya

[‡]University of Belgrade - Institute for Biological Research "Sinisa Stankovic," Bulevar Despota Stefana 142, 11000 Belgrade, Serbia

[§]University of Belgrade - Faculty of Chemistry, Studentski trg 12-16, 11000 Belgrade, Serbia

ABSTRACT

The essential oil of *Thymus algeriensis* was analyzed as a potential preservative in soft cheese. We developed a novel method to test the preserving properties of essential oil in soft cheese. Contamination incidence of *Penicillium aurantiogriseum* was absent after 30 d of storage at 4°C with 25 µL of essential oil added. The antimicrobial activity was tested against 8 bacteria and 8 fungi. *Thymus algeriensis* oil showed inhibitory activity against tested bacteria at 0.03 to 0.09 mg/mL, and bactericidal activity was achieved at 0.05 to 0.15 mg/mL. For antifungal activity, minimum inhibitory concentrations ranged between 0.01 and 0.04 mg/mL and minimum fungicidal concentrations between 0.01 and 0.04 mg/mL. Furthermore, the oil was also screened for antiradical activity using the 1,1-diphenyl-2-picrylhydrazyl assay. The results showed that the oil was active and achieved half-maximal inhibitory activity at 0.132 mg/mL. We used gas chromatography, gas chromatography-mass spectrometry, and nuclear magnetic resonance spectrometry to investigate the volatile compounds from the oil. Carvacrol was identified as the main compound in the oil, represented by 80.9% of the total constituents, followed by *p*-cymene (7.7%).

Key words: *Thymus algeriensis*, essential oil, carvacrol, antimicrobial, cheese preservation

Short Communication

Dairy products can be easily contaminated by pathogens or food spoilage microorganisms, which can produce unwanted appearance, decrease commercial value, and, most importantly, compromise the safety

of products. Molds on the surface of cheese cause undesirable flavors and many quality problems. The most frequently isolated mold genus from cheese is *Penicillium* spp., which may produce mycotoxins such as ochratoxin A and citrinin (Torkar and Vengust, 2008). Some studies have shown that natural compounds may protect cheese from food contaminant microorganisms (Smith-Palmer et al., 2001; Lucera et al., 2012); Gammariello et al. (2010) reported that chitosan and lemon extract improve shelf life of Mediterranean fresh cheese.

Plant products and their derived compounds are of growing interest due to the wide range of bioactivities that make them useful as natural additives in different types of food. For centuries, they have been used for treatment of infections and diseases worldwide (Rios et al., 1988; Skrinjar and Nemet, 2009; Gottardi et al., 2016). In recent years, utilization of plant products [e.g., essential oils (EO), extracts, and their secondary metabolites] has gained considerable interest for the food industry as an alternative to synthetic antimicrobials or additives (Bukvicki et al., 2014a,b, 2015; Tyagi et al., 2014).

The majority of the thyme species are widely used as medicinal herbs (Oussalah et al., 2007) and known to have biological activities, such as antispasmodic (Babaei et al., 2008), sedative (Fachini-Queiroz et al., 2012), antioxidant (Lee et al., 2005), and antimicrobial (Oussalah et al., 2007; Hazzit et al., 2009) activities. The focus of our study was potential application of *Thymus algeriensis* EO in soft cheese, and we aimed to investigate the effect of the oil as preservative in this dairy product. For such purposes, we developed a novel method to test antimicrobial food-preserving properties of substances in soft cheese. Furthermore, chemical characterization of the EO and its antiradical attributes and antimicrobial activities were investigated.

Thyme plants were collected from Gharyan city (Al Gabel Al-Garbe, Libya) in April 2012. The plant was identified by A. Giweli and then confirmed by P. D. Marin. One hundred grams of air-dried aerial parts of

Received August 19, 2017.

Accepted December 12, 2017.

¹Corresponding author: dankabukvicki@bio.bg.ac.rs, dankabukvicki@gmail.com

plant samples, with woody parts removed, were submitted to hydrodistillation (Bukvicki et al., 2014a).

The volatile composition of the *T. algeriensis* EO was determined according to a previously published procedure (Bukvicki et al., 2016). The components were identified by comparison of mass spectra with reference from commercial libraries (Wiley 7, NIST 11, and Adams 4; McLafferty, 2005; Adams, 2007; NIST, 2011). Structural determination of each component was confirmed by comparison of retention indices (RI) obtained by the standard method (Vandendool and Kratz, 1963) with the Adams 4 RI library. Quantitative ^1H nuclear magnetic resonance (NMR) analysis of the *T. algeriensis* EO was performed by using a Bruker Avance III 500 NMR spectrometer (500 MHz, Bruker, Rheinstetten, Germany) in deuterated chloroform (CDCl_3), with tetramethylsilane as a reference and 2,6-bis(1,1-dimethylethyl)-4-methylphenol (butylated hydroxytoluene, **BHT**) as an internal standard. Essential oil (5.260 mg) and BHT (5.050 mg) were dissolved in 500 μL of CDCl_3 in a 5-mm NMR tube. Proton NMR spectra were run with a standard pulse sequence with 16 scans, and with 3 different relaxation delays (d1) times of 1, 4, and 10 s. The BHT was selected because its ^1H NMR signals (aromatic proton) are clearly distinguishable from those of EO components (Figure 1). Peak area of characteristic protons used for quantitative calculation was manually integrated. Signal of 1 proton of carvacrol [chemical shift (δ) 7.03 doublet, coupling constant (J) = 7.7 Hz], 2 signals of 4 protons of *p*-cymene (δ 7.07–7.15 multiplets), and signal of 2 protons of BHT (δ 6.98 singlet) were used for relative percentage calculation, based on the equation

$$Px(\%) = \frac{Mx}{Ms} \times \frac{m_s}{m_{EO}} \times \frac{Ix}{Is} \times \frac{n_s}{n_x} \times 100,$$

where Mx and Ms are the molar masses of the analyte and the internal standard (BHT); m_{EO} and m_s are the weighed mass of the essential oil of *T. algeriensis* and BHT; Ix and Is are the peak areas of the analyte and BHT; and n_x and n_s are the numbers of the protons generating the corresponding peak of the analyte and BHT, respectively. The 1,1-diphenyl-2-picrylhydrazyl (**DPPH**) assay was done according to a previously published procedure (Dzamić et al., 2015), as were the assays for antibacterial and antifungal activity (Bukvicki et al., 2016).

Soft cheese used in our experiment was obtained from a market located in Belgrade (Serbia). Soft cheese (domaći mladi sir, without preservatives; Président/Group Lactalis, Laval, France), with nutritional values of 5.5% milk fat, 4% carbohydrates, and 12.5% protein,

was stored at 4°C for until further use. Cheese was sliced into equal cuboid shapes, with an approximate weight per slice of 20 g, and placed in sterile petri dishes. For the in situ antifungal assay, we developed a convenient assay for all investigations. Slices of soft cheese were pierced (5 mm deep and 5 mm wide), creating 3 wells per slice. Each treatment had 2 replicates with 5 slices of cheese per replicate. Essential oil of *T. algeriensis* was dissolved in different concentrations of sterile 0.2% Tween 80 (vol/vol). Essential oils were applied at 1, 5, 10, 15, 20, and 25 μL , and then 25 μL of EO emulsion was placed into each well made in cheese. After 1 h, 25 μL of a conidial suspension (2×10^4 conidia/mL) of *Penicillium aurantiogriseum* was added to each well in sliced cheese. As a positive control of fungal growth, *P. aurantiogriseum* was inoculated on cheese without EO. The treated slices of cheese were stored at 4°C for 30 d, respectively. After storage, the number of wells that showed contamination symptoms (fungal growth) was recorded, and contamination incidence (%) was calculated as

$$\text{Contamination incidence} = \left(\frac{\text{number of infected wells}}{\text{total wells per replicate}} \right) \times 100.$$

To evaluate possible changes in color, texture, and flavor, cheese was sprayed with EO at levels of 1, 5, and 15 μL per surface of sliced cheese [diluted EO at 1, 5, and 15 μL in 100 μL of 0.2% Tween 80 (vol/vol) in water], with 1 control sample not sprayed with EO. Test was made in initial shelf-life, third day after production. Results were expressed as average grades given by 20 panelists (Bukvicki et al., 2014a).

Based on GC and GC-MS analysis, 21 compounds were identified, representing 99.7% of the total oil. Carvacrol was the main compound in the oil (80.9% of total oil), followed by *p*-cymene (7.7%). In addition, 2 components of analyzed EO were identified and quantified by ^1H NMR (Figure 1). Relative mass percentage of carvacrol was 72.8 and *p*-cymene 7.6; moreover, substitutional isomers of cymene (*o*-, *m*-, *p*-) have similar mass spectra, and RI, GC, and GC-MS analysis could be time consuming. Otherwise, differences among proton NMR spectra of isomers of cymene provide clear identification. The NMR analysis could be the method for quality control of chemical composition of EO with few dominant compounds.

Thymus algeriensis EO, its main component carvacrol, and butylated hydroxyl anisole had good anti-radical potential as investigated by the DPPH assay (results in Figure 2). Our investigations showed that *T. algeriensis* EO exhibited notable free radical scavenging activity, with half-maximal inhibitory concentra-

tion (IC_{50}) of 0.132 mg/mL, although it was weaker than carvacrol alone ($IC_{50} = 0.105$ mg/mL). The good antiradical potential of the oil could be attributed to carvacrol, as such a relation was previously shown for oxygenated monoterpenes by Hazzit et al. (2009).

The results of the antimicrobial activity tested by microdilution method are summarized in Table 1. The EO of *T. algeriensis* exhibited significant antimicrobial activity against all strains tested. For antibacterial activity, the inhibition values ranged from an MIC of 0.03 to 0.09 mg/mL and minimum bacterial concentration of 0.05 to 0.18 mg/mL. It can be seen that the oil showed activity better than the positive control (ampicillin) and streptomycin, which showed the lowest activity. For antifungal activity, the oil was active at 0.01 to 0.04 mg/mL and minimum fungicidal concentration (MFC) of 0.01 to 0.04 mg/mL, whereas the control ketoconazole achieved MIC at 0.15 to 1.5 mg/mL and MFC at 0.2 to 2.0 mg/mL, and bifonazole showed MIC

at 0.1 to 0.2 mg/mL and MFC at 0.2 to 0.3 mg/mL (Table 2, Table 3).

Generally, the fungi appear to be more sensitive than the bacteria. Correlation between the achieved antimicrobial activity of selected EO and their chemical composition leads us to infer that the activity may be ascribed to carvacrol, which was present at a high percentage in the oils. Carvacrol has been shown to be a good antimicrobial agent in previous studies (Nostro and Papalia, 2012). Previous reports showed that carvacrol inhibits growth of *Aspergillus niger* at 50 μ g/mL, *Aspergillus fumigates*, *Micrococcus flavus*, and *Aspergillus ochraceus* at 100 μ g/mL, as well as *Penicillium chrysogenum* at 500 μ g/mL (Abbaszadeh et al., 2014). The MIC of carvacrol against 16 *Candida albicans* strains was 256 μ g/mL and MFC was 512 μ g/mL (Lima et al., 2013). Against 4 *Streptococcus* strains, carvacrol showed MIC of 2.5 mg/mL and minimum bacterial concentration of 5 mg/mL, whereas against

Figure 1. Quantitative 1H nuclear magnetic resonance (500 MHz, deuterated chloroform) analysis of the essential oil of *Thymus algeriensis* with 2,6-bis(1,1-dimethylethyl)-4-methylphenol (BHT) as an internal standard. Expansion showed signals from aromatic protons of BHT, carvacrol, and *p*-cymene used for quantitative calculations.

Table 1. Chemical composition of *Thymus algeriensis* essential oil analyzed by GC-MS and GC-flame-ionization detection¹

t_R (min)	RI	Compound	%
5.52	922	α -Thujene	0.2
5.69	929	α -Pinene	0.5
6.09	943	Camphene	0.1
6.88	972	β -Pinene	<0.1%
7.26	986	Myrcene	0.9
7.72	1,002	α -Phellandrene	0.1
7.90	1,007	3-Carene	0.1
8.10	1,012	α -Terpinene	0.8
8.39	1,020	<i>p</i> -Cymene	7.7
8.54	1,024	β -Phellandrene	0.5
9.60	1,054	γ -Terpinene	3.6
10.76	1,086	Terpinolene	<0.1%
11.21	1,098	Linalool	1.1
13.95	1,161	Borneol	0.5
14.49	1,174	Terpinene-4-ol	0.6
15.21	1,190	α -Terpineol	0.1
19.61	1,290	Thymol	0.3
20.23	1,300	Carvacrol	80.9
25.07	1,415	β -Caryophyllene	1.2
26.54	1,450	α -Humulene	<0.1%
31.95	1,580	Caryophyllene oxide	0.5
		Total	99.7

¹ t_R = retention time on HP-5MSI capillary column (30 m $\times$ 0.25 mm i.d., film thickness 0.25 μ m,) RI = retention indices computed by comparison with retention times of *n*-alkanes under the same chromatographic conditions (Vandendool and Kratz, 1963); relative percentage calculated by GC-FID peak areas.

Candida albicans MIC was 1.25 mg/mL and MFC was 2.5 mg/mL (Botelho et al., 2007). Ramos et al. (2013) showed that carvacrol is safe for use with food for human consumption.

The results of in situ treatment of sliced cheese with EO of *T. algeriensis* are given in Table 4 as contamination incidence (%). For 30 d after treatment and inoculation with *P. aurantiogriseum*, pieces of sliced cheese were kept at 4°C. The incidence of contamination decreased with increasing EO concentration, with contamination being completely absent at the highest dose used. Results showed that 100% of control sliced cheese initially developed infection (Figure 3). Contamination incidence of *P. aurantiogriseum* in sliced cheese decreased from 66 to 0.0% with increasing concentrations of EO from 20 to 25 μ L/mL, respectively. Lower concentrations of EO had no effect on contamination incidence and it remained at 100%. Higher concentrations of EO needed to prevent cheese contamination and the low contamination incidence, in contrast to the in vitro microdilution method, suggested that cheese is a more complex medium than the laboratory broths used in in vitro antimicrobial assays. In commonly used doses (up to 20 drops per day; as a culinary herb), no acute or chronic toxicity has been reported for thyme

Figure 2. Half-maximal inhibitory concentration (IC₅₀) values with standard deviations of essential oil tested in a 1,1-diphenyl-2-picrylhydrazyl assay with its principal compound carvacrol and positive control butylated hydroxyl anisole (BHA). Color version available online.

Table 2. Antibacterial activity (mg/mL) of *Thymus algeriensis* essential oil against food spoilage bacteria

Agent	<i>Staphylococcus aureus</i> (ATCC 6538)	<i>Salmonella</i> Typhimurium (ATCC 13311)	<i>Enterobacter cloacae</i> (human isolate)	<i>Escherichia coli</i> (ATCC 35210)	<i>Pseudomonas aeruginosa</i> (ATCC 27853)	<i>Listeria monocytogenes</i> (NCTC 7973)	<i>Micrococcus flavus</i> (ATCC 10240)	<i>Bacillus cereus</i> (clinical isolate)
<i>T. algeriensis</i>								
MIC	0.08 ± 0.03 ^b	0.09 ± 0.04 ^a	0.05 ± 0.04 ^a	0.08 ± 0.03 ^a	0.05 ± 0.00 ^a	0.04 ± 0.00 ^a	0.03 ± 0.00 ^a	0.04 ± 0.01 ^a
MBC	0.15 ± 0.05 ^b	0.18 ± 0.07 ^a	0.11 ± 0.07 ^a	0.11 ± 0.07 ^a	0.11 ± 0.01 ^a	0.09 ± 0.02 ^a	0.05 ± 0.00 ^a	0.08 ± 0.02 ^a
Ampicillin								
MIC	0.05 ± 0.03 ^a	0.20 ± 0.00 ^b	0.30 ± 0.02 ^b	0.20 ± 0.03 ^b	0.20 ± 0.02 ^b	0.20 ± 0.02 ^b	0.20 ± 0.01 ^b	0.10 ± 0.03 ^b
MBC	0.10 ± 0.03 ^a	0.30 ± 0.02 ^b	0.50 ± 0.02 ^b	0.300 ± 0.006 ^b	0.30 ± 0.02 ^b	0.30 ± 0.09 ^b	0.30 ± 0.02 ^b	0.20 ± 0.01 ^b
Streptomycin								
MIC	0.300 ± 0.003 ^c	0.30 ± 0.01 ^c	0.40 ± 0.02 ^c	0.300 ± 0.006 ^c	0.80 ± 0.01 ^c	0.400 ± 0.006 ^c	0.30 ± 0.01 ^c	0.30 ± 0.00 ^c
MBC	0.400 ± 0.006 ^c	0.50 ± 0.02 ^c	0.80 ± 0.01 ^c	0.500 ± 0.006 ^c	1.25 ± 0.02 ^c	0.50 ± 0.01 ^c	0.40 ± 0.02 ^c	0.40 ± 0.01 ^c

^{a-c}In each column, and for MIC or minimum bactericidal concentration values, different letters mean significant differences between *T. algeriensis* essential oil, ampicillin, and streptomycin ($P < 0.05$).

Table 3. Antifungal activity (mg/mL) of *Thymus algeriensis* essential oil and antifungal agents against food spoilage fungi (*Aspergillus*, *Trichoderma*, and *Penicillium* spp.)

Agent	<i>A. fumigates</i> (human isolate)	<i>A. versicolor</i> (ATCC 11730)	<i>A. ochraceus</i> (ATCC 12066)	<i>A. niger</i> (ATCC 6275)	<i>Tr. viride</i> (IAM 5061)	<i>P. funiculosum</i> (ATCC 36839)	<i>P. ochrochloron</i> (ATCC 9112)	<i>P. aurantiogriseum</i> (food isolate)
<i>T. algeriensis</i>								
MIC	0.01 ± 0.00 ^a	0.04 ± 0.03 ^a	0.01 ± 0.00 ^a	0.01 ± 0.00 ^a	0.01 ± 0.00 ^a	0.01 ± 0.01 ^a	0.01 ± 0.02 ^a	0.02 ± 0.01 ^a
MFC	0.03 ± 0.00 ^a	0.03 ± 0.01 ^a	0.03 ± 0.00 ^a	0.01 ± 0.00 ^a	0.01 ± 0.00 ^a	0.03 ± 0.02 ^a	0.03 ± 0.02 ^a	0.04 ± 0.01 ^a
Ketoconazole								
MIC	0.20 ± 0.01 ^c	0.20 ± 0.06 ^c	0.15 ± 0.00 ^b	0.20 ± 0.03 ^c	1.00 ± 0.00 ^c	0.20 ± 0.03 ^b	1.00 ± 0.30 ^c	1.50 ± 0.00 ^c
MFC	0.50 ± 0.06 ^c	0.50 ± 0.06 ^c	0.20 ± 0.30 ^b	0.50 ± 0.03 ^c	1.50 ± 0.30 ^c	0.50 ± 0.00 ^c	1.50 ± 0.10 ^c	2.00 ± 0.00 ^c
Bifonazole								
MIC	0.15 ± 0.03 ^b	0.10 ± 0.06 ^b	0.15 ± 0.03 ^b	0.15 ± 0.03 ^b	0.15 ± 0.03 ^b	0.20 ± 0.06 ^b	0.20 ± 0.06 ^b	0.20 ± 0.00 ^b
MFC	0.20 ± 0.06 ^b	0.20 ± 0.06 ^b	0.20 ± 0.00 ^b	0.20 ± 0.06 ^b	0.20 ± 0.00 ^b	0.25 ± 0.03 ^b	0.25 ± 0.03 ^b	0.30 ± 0.00 ^b

^{a-c}In each column, and for MIC or minimum fungicidal concentration values, different letters indicate significant differences between *T. algeriensis* EO, ketoconazole, and bifonazole ($P < 0.05$).

Figure 3. Contamination of sliced cheese with *Penicillium aurantiogriseum* after 30 d of storage at 4°C. (A) Control sample not treated with essential oil of *Thymus algeriensis* showing fully developed contamination; (B) sample of cheese treated with concentration of 1 µL per well in cheese; and (C) sample of cheese treated with the highest concentration of 15 µL per well in cheese, showing no contamination. Color version available online.

oil (Mills and Bone, 2000); thus, the applied doses in our study are considered as safe for use.

Results of sensory evaluation of cheese sprayed with EO showed no changes in texture and color on surface of cheese slices after 30 d of storage at temperature 4°C (Table 5). The results indicated increasing grades

Table 4. Incidence of contamination with *Penicillium aurantiogriseum* of cheese treated with essential oil of *Thymus algeriensis* (EO)

EO (µL/well)	Contamination incidence (%)
0	100.0
1	100.0
5	100.0
10	100.0
15	100.0
20	66.0
25	0.0

Table 5. Changes in color, texture, and taste during storage of 30 d at 4°C¹ in control cheese slices and samples with different concentrations of essential oils (EO) added

Item	Control (no EO)	EO added per well in cheese (µL)		
		1	5	15
Color	3.6	4.4	4.5	4.6
Texture	4.2	3.9	4.0	4.2
Taste	4.3	3.7	3.6	3.1

¹The results are expressed as the average of all grades: 1 = extremely dislike, 2 = dislike, 3 = neither like nor dislike, 4 = like; 5 = extremely like.

of color with addition of EO, whereas cheese taste was decreased but still was above the limits for nonliked products.

Thymus algeriensis EO showed antimicrobial potential against the food pathogenic mold *P. aurantiogriseum* in vitro as well as in real food (cheese) systems. The high amounts of carvacrol present in the oil are likely responsible for the antimicrobial activity. *Thymus* species are well-known culinary spices widely used all around the world, with pleasant taste and flavor. With high antifungal and antiradical abilities, this EO could be recommended as natural antimicrobial additive for shelf-life prolongation of the soft cheese.

ACKNOWLEDGMENTS

This research was supported by a grant from the Ministry of Education, Science and Technological Development of Serbia (Projects No. 173029, 173032 and 172053).

REFERENCES

Abbaszadeh, S., A. Sharifzadeh, H. Shokri, A. R. Khosravi, and A. Abbaszadeh. 2014. Antifungal efficacy of thymol, carvacrol, eugenol and menthol as alternative agents to control the growth of food-relevant fungi. *J. Mycol. Med.* 24:e51–56.

Adams, R. P. 2007. Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry. 4th ed. Allured Publ., Carol Stream, IL.

Babaei, M., M. E. Abarghoei, R. Ansari, A. A. Vafaei, A. A. Taherian, M. M. Akhavan, G. Toussy, and S. Mousavi. 2008. Antispasmodic effect of hydroalcoholic extract of *Thymus vulgaris* on the guinea pig-ileum. *Nat. Prod. Res.* 22:1143–1150.

- Botelho, M. A., N. A. P. Nogueira, G. M. Bastos, S. G. C. Fonseca, T. L. G. Lemos, F. J. A. Matos, D. Montenegro, J. Heukelbach, V. S. Rao, and G. A. C. Brito. 2007. Antimicrobial activity of the essential oil from *Lippia sidoides*, carvacrol and thymol against oral pathogens. *Braz. J. Med. Biol. Res.* 40:349–356.
- Bukvicki, D., A. Cirić, M. Soković, L. Vannini, L. Nissen, M. Novaković, Lj. Vujisić, Y. Asakawa, and P. D. Marin. 2016. *Micromeria thymifolia* essential oil suppresses quorum-sensing signaling in *Pseudomonas aeruginosa*. *Nat. Prod. Commun.* 11:1903–1906.
- Bukvicki, D., D. Gottardi, A. K. Tyagi, M. Veljic, P. D. Marin, Lj. Vujisić, M. E. Guerzoni, and L. Vannini. 2014b. *Scapania nemorea* liverwort extracts: Investigation on volatile compounds, in vitro antimicrobial activity and control of *Saccharomyces cerevisiae* in fruit juice. *Lebensm. Wiss. Technol.* 55:452–458.
- Bukvicki, D., D. Stojkovic, M. Sokovic, M. Nikolic, L. Vannini, C. Montanari, and P. D. Marin. 2015. Potential application of *Micromeria dalmatica* essential oil as a protective agent in a food system. *Lebensm. Wiss. Technol.* 63:262–267.
- Bukvicki, D., D. Stojkovic, M. Sokovic, L. Vannini, C. Montanari, B. Pejin, A. Savic, M. Veljic, S. Grujic, and P. D. Marin. 2014a. *Satureja horvatii* essential oil: In vitro antimicrobial and antiradical properties and in situ control of *Listeria monocytogenes* in pork meat. *Meat Sci.* 96:1355–1360.
- Dzamić, A., B. Nikolic, A. Giweli, D. Mitic-Culafic, M. Sokovic, M. Ristic, J. Knezevic-Vukcevic, and P. Marin. 2015. Libyan *Thymus capitatus* essential oil: Antioxidant, antimicrobial, cytotoxic and colon pathogen adhesion-inhibition properties. *J. Appl. Microbiol.* 119:389–399.
- Fachini-Queiroz, F.C., R. Kummer, C.F. Estevao-Silva, M.D. Carvalho, J. M. Cunha, R. Grespan, C. A Bersani-Amado, and R. K. Nakamura Cuman. 2012. Effects of thymol and carvacrol, constituents of *Thymus vulgaris* L. essential oil, on the inflammatory response. *Evid. Based Complement. Altern. Med.* 2012:657026. <https://doi.org/10.1155/2012/657026>.
- Gammariello, D., A. Conte, and M. A. Del Nobile. 2010. Assessment of chitosan and extracts of lemon and sage as natural antimicrobial agents during Fior di latte cheese-making. *Int. J. Dairy Technol.* 63:530–537.
- Gottardi, D., D. Bukvicki, S. Prasad, and A. K. Tyagi. 2016. Beneficial effects of spices in food preservation and safety. *Front. Microbiol.* 7:1394.
- Hazzit, M., A. Baaliouamer, A. R. Verissimo, M. L. Faleiro, and M. G. Miguel. 2009. Chemical composition and biological activities of Algerian *Thymus* oils. *Food Chem.* 116:714–721.
- Lee, S. J., K. Umamo, T. Shibamoto, and K. G. Lee. 2005. Identification of volatile components in basil (*Ocimum basilicum* L.) and thyme leaves (*Thymus vulgaris* L.) and their antioxidant properties. *Food Chem.* 91:131.
- Lima, I. O., F. O. de Oliveira Pereira, W. A. de Oliveira, E. de Oliveira Lima, E. A. Menezes, F. A. Cunha, and M. F. F. M. Diniz. 2013. Antifungal activity and mode of action of carvacrol against *Candida albicans* strains. *J. Essent. Oil Res.* 25:138–142. <https://doi.org/10.1080/10412905.2012.754728>.
- Lucera, A., C. Costa, A. Conte, and M. A. Del Nobile. 2012. Food applications of natural antimicrobial compounds. *Front. Microbiol.* 3:287.
- McLafferty, F. W. 2005. Wiley Registry of Mass Spectral Data: With NIST 2005 Spectral Data, 7th rev. ed., John Wiley and Sons Ltd., New York, NY.
- Mills, S., and K. Bone. 2000. Principles and Practice of Phytotherapy. Churchill Livingstone, Toronto, Canada.
- NIST. 2011. NIST/NIH/EPA Mass Spectral Library, Standard Reference Database 1, NIST 11. Standard Reference Data Program, National Institute of Standards and Technology, Gaithersburg, MD.
- Nostro, A., and T. Papalia. 2012. Antimicrobial activity of carvacrol: current progress and future perspectives. *Recent Pat. Antiinfect. Drug Discov.* 7:28–35.
- Oussalah, M., S. Caille, L. Saucier, and M. Lacroix. 2007. Inhibitory effects of selected plant essential oils on the growth of four pathogenic bacteria: *E. coli* O157:H7, *Salmonella typhimurium*, *Staphylococcus aureus* and *Listeria monocytogenes*. *Food Control* 18:414–420.
- Ramos, M., A. Beltran, A. Valdes, M. A. Peltzer, A. Jimenez, M. A. Peltzer, A. Jimenez, M. C. Garrigos, and G. E. Zaikov. 2013. Carvacrol and Thymol for Fresh Food Packaging. *J. Bioequiv.* 5:154–160. <https://doi.org/10.4172/jbb.1000151>.
- Rios, J. L., M. C. Recio, and A. Villar. 1988. Screening methods for natural antimicrobial products with antimicrobial activity: A review of the literature. *J. Ethnopharmacol.* 23:127–149.
- Skrinjar, M., and N. Nemet. 2009. Antimicrobial effects of spices and herbs essential oils. *Acta Period. Technol.* 40:195–209.
- Smith-Palmer, A., J. Stewart, and L. Fyfe. 2001. The potential application of plant essential oils as natural food preservatives in soft cheese. *Food Microbiol.* 18:463–470.
- Torkar, K. G., and A. Vengust. 2008. The presence of yeasts, moulds and aflatoxin M1 in raw milk and cheese in Slovenia. *Food Control* 19:570–577.
- Tyagi, A. K., D. Bukvicki, D. Gottardi, G. Tabanelli, C. Montanari, A. Malik, and M. E. Guerzoni. 2014. Eucalyptus essential oil as a natural food preservative: In vivo and in vitro antiyeast potential. *BioMed Res. Int.* 2014:969143. <https://doi.org/10.1155/2014/969143>.
- Vandendool, H., and P. D. Kratz. 1963. A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *J. Chromatogr.* 11:463–471.