Corrigendum to “Addition of straw to the early-lactation diet: Effects on feed intake, milk yield, and subclinical ketosis in Holstein cows” (J. Dairy Sci. 104:3008–3017)

Hesam A. Seifi, Julianna M. Huzzey, M. A. Khan, Daniel M. Weary, and Marina A. G. von Keyserlingk

In Table 1 (page 3010), the authors presented the calculated ration formulation and not the analytical values of the ration composition. The revised table is shown below with corrected values in bold.

The authors regret the error.

REFERENCES

Seifi, H. A., J. M. Huzzey, M. A. Khan, D. M. Weary, and M. A. G. von Keyserlingk. 2021. Addition of straw to the early-lactation diet: Effects on feed intake, milk yield, and subclinical ketosis in Holstein cows. J. Dairy Sci. 104:3008–3017. <https://doi.org/10.3168/jds.2020-18549>.

Table 1. Composition and component analysis (% of DM unless otherwise noted) of composite samples of 3 TMR types fed to transition cows: dry-cow TMR (DTMR), conventional postpartum TMR (CTMR), and postpartum TMR with added wheat straw (STMR; fed only during the 3-wk postcalving treatment period)¹

Composition	DTMR	CTMR	STMR
Ingredient			
Grass silage	13.07	22.25	21.00
Corn silage	30.25	16.64	15.93
Alfalfa hay	16.43	11.72	11.22
Concentrate mix ²	14.17	49.38	47.27
Wheat straw	26.08	0.00	4.27
Component			
DM	49.28	52.45	53.41
NE _L (Mcal/kg)	1.45	1.57	1.57
TDN	63.75	68.63	68.67
CP	13.56	16.09	16.58
NDF	44.34	37.55	38.11
ADF	32.29	26.02	25.97
NFC	30.60	34.86	33.80
Starch	11.57	15.53	16.85
Crude fat	2.18	2.93	2.89
Ash	8.34	9.50	9.24
Ca	0.94	1.05	0.91
P	0.26	0.33	0.34
Mg	0.32	0.31	0.30
Na	0.16	0.38	0.37
K	1.57	1.78	1.80
S	0.19	0.22	0.21

¹All cows were fed the CTMR after the treatment period (n = 34 cows per treatment).

²For DTMR, the major ingredients included 45% Amino Plus (Ag Processing Inc., Omaha, NE), 12% corn gluten meal, 11% soybean meal, 7.36% fine limestone, 5% ground beet pulp, 5% dry cow mineral, 4.32% calcium sulfate, 3.544% distillers corn wheat blend, 1% molasses. CTMR and STMR: 34% rolled barley, 20% fine ground barley, 10% distillers corn wheat blend, 7.3% ground wheat, 6% soybean meal, 6% Amino Plus (Ag Processing Inc.), 5% canola meal, 2.22% fine limestone, 1.18% mill run pellets. (Analysis conducted by A&L Canada Laboratories Inc., London, ON, Canada.)